

A sharp Trudinger-Moser type inequality for unbounded domains in $\mathbb{R}^n$

Yuxiang Li and Bernhard Ruf

Abstract

The Trudinger-Moser inequality states that for functions $u \in H_0^{1,n}(\Omega)$ ($\Omega \subset \mathbb{R}^n$ a bounded domain) with $\int_{\Omega} |\nabla u|^n dx \leq 1$ one has $\int_{\Omega} (e^{\alpha_n |u|^{\frac{n}{n-1}}} - 1) dx \leq c |\Omega|$, with c independent of u . Recently, the second author has shown that for $n = 2$ the bound $c |\Omega|$ may be replaced by a uniform constant d independent of Ω if the Dirichlet norm is replaced by the Sobolev norm, i.e. requiring $\int_{\Omega} (|\nabla u|^n + |u|^n) dx \leq 1$. We extend here this result to arbitrary dimensions $n > 2$. Also, we prove that for $\Omega = \mathbb{R}^n$ the supremum of $\int_{\mathbb{R}^n} (e^{\alpha_n |u|^{\frac{n}{n-1}}} - 1) dx$ over all such functions is attained. The proof is based on a blow-up procedure.

Keywords: Trudinger-Moser inequality, blow-up, best constant, unbounded domain.

Mathematics subject classification (2000): 35J50, 46E35

1 Introduction

Let $H_0^{1,p}(\Omega)$, $\Omega \subseteq \mathbb{R}^n$, be the usual Sobolev space, i.e. the completion of $C_0^\infty(\Omega)$ with the norm

$$\|u\|_{H^{1,p}(\Omega)} = \left(\int_{\Omega} (|\nabla u|^p + |u|^p) dx \right)^{\frac{1}{p}}.$$

It is well-known that

$$\begin{aligned} H_0^{1,p}(\Omega) &\subset L^{\frac{pn}{n-p}}(\Omega) \quad \text{if } 1 \leq p < n \\ H_0^{1,p}(\Omega) &\subset L^\infty(\Omega) \quad \text{if } n < p \end{aligned}$$

The case $p = n$ is the limit case of these embeddings and it is known that

$$H_0^{1,n}(\Omega) \subset L^q(\Omega) \quad \text{for } n \leq q < +\infty.$$

When Ω is a bounded domain, we usually use the Dirichlet norm $\|u\|_D = (\int |\nabla u|^n dx)^{\frac{1}{n}}$ in place of $\|\cdot\|_{H^{1,n}}$. In this case, we have the famous Trudinger-Moser inequality (see [P], [T], [M]) for the limit case $p = n$ which states that

$$\sup_{\|u\|_D \leq 1} \int_{\Omega} (e^{\alpha |u|^{\frac{n}{n-1}}} - 1) dx = c(\Omega, \alpha) \begin{cases} < +\infty & \text{when } \alpha \leq \alpha_n \\ = +\infty & \text{when } \alpha > \alpha_n \end{cases} \quad (1.1)$$

where $\alpha_n = n\omega_{n-1}^{\frac{1}{n-1}}$, and ω_{n-1} is the measure of the unit sphere in $\mathbb{R}^n$. The Trudinger-Moser result has been extended to Sobolev spaces of higher order and Sobolev spaces over compact

manifolds (see [A], [Fo]). Moreover, for any bounded Ω , the constant $c(\Omega, \alpha_n)$ can be attained. For the attainability, we refer to [C-C], [F], [Lin], [L1], [L2], [d-d-R], [L3].

Another interesting extension of (1.1) is to construct Trudinger-Moser type inequalities on unbounded domains. When $n = 2$, this has been done by B. Ruf in [R]. On the other hand, for an unbounded domain in $\mathbb{R}^n$, S. Adachi and K. Tanaka ([A-T]) get a weaker result. Let

$$\Phi(t) = e^t - \sum_{j=1}^{n-2} \frac{t^j}{j!}.$$

S. Adachi and K. Tanaka's result says that:

Theorem A *For any $\alpha \in (0, \alpha_n)$ there is a constant $C(\alpha)$ such that*

$$\int_{\mathbb{R}^n} \Phi\left(\alpha \left(\frac{|u|}{\|\nabla u\|_{L^n(\mathbb{R}^n)}}\right)^{\frac{n}{n-1}}\right) dx \leq C(\alpha) \frac{\|u\|_{L^n(\mathbb{R}^n)}^n}{\|\nabla u\|_{L^n(\mathbb{R}^n)}^n}, \quad \text{for } u \in H^{1,n}(\mathbb{R}^n) \setminus \{0\}. \quad (1.2)$$

In this paper, we shall discuss the critical case $\alpha = \alpha_n$. More precisely, we prove the following:

Theorem 1.1. *There exists a constant $d > 0$, s.t. for any domain $\Omega \subset \mathbb{R}^n$,*

$$\sup_{u \in H^{1,n}(\Omega), \|u\|_{H^{1,n}(\Omega)} \leq 1} \int_{\Omega} \Phi(\alpha_n |u|^{\frac{n}{n-1}}) dx \leq d. \quad (1.3)$$

The inequality is sharp: for any $\alpha > \alpha_n$, the supremum is $+\infty$.

We set

$$S = \sup_{u \in H^{1,n}(\mathbb{R}^n), \|u\|_{H^{1,n}(\mathbb{R}^n)} \leq 1} \int_{\mathbb{R}^n} \Phi(\alpha_n |u|^{\frac{n}{n-1}}) dx.$$

Further, we will prove

Theorem 1.2. *S is attained. In other words, we can find a function $u \in H^{1,n}(\mathbb{R}^n)$, with $\|u\|_{H^{1,n}(\mathbb{R}^n)} = 1$, s.t.*

$$S = \int_{\mathbb{R}^n} \Phi(\alpha_n |u|^{\frac{n}{n-1}}) dx.$$

The second part of Theorem 1.1 is trivial: Given any fixed $\alpha > \alpha_n$, we take $\beta \in (\alpha_n, \alpha)$. By (1.1) we can find a positive sequence $\{u_k\}$ in

$$\{u \in H_0^{1,n}(B_1) : \int_{B_1} |\nabla u|^n dx = 1\},$$

such that

$$\lim_{k \rightarrow +\infty} \int_{B_1} e^{\beta u_k^{\frac{n}{n-1}}} = +\infty.$$

By Lion's Lemma, we get $u_k \rightarrow 0$. Then by the compact embedding theorem, we may assume $\|u_k\|_{L^p(B_1)} \rightarrow 0$ for any $p > 1$. Then, $\int_{\mathbb{R}^n} (|\nabla u_k|^n + |u_k|^n) dx \rightarrow 1$, and

$$\alpha \left(\frac{u_k}{\|u_k\|_{H^{1,n}}}\right)^{\frac{n}{n-1}} > \beta u_k^{\frac{n}{n-1}}$$

when k are sufficiently large. So, we get

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\alpha(\frac{u_k}{\|u_k\|_{H^{1,n}}})^{\frac{n}{n-1}}) dx \geq \lim_{k \rightarrow +\infty} \int_{B_1} (e^{\beta u_k^{\frac{n}{n-1}}} - 1) dx = +\infty.$$

The first part of Theorem 1.1 and Theorem 1.2 will be proved by blow up analysis. We will use the ideas from [L1] and [L2] (see also [A-M] and [A-D]). However, in the unbounded case we do not obtain the strong convergence of u_k in $L^n(\mathbb{R}^n)$, and so we need more techniques.

Concretely, we will find positive and symmetric functions $u_k \in H_0^{1,n}(B_{R_k})$ which satisfy

$$\int_{B_{R_k}} (|\nabla u_k|^n + |u_k|^n) dx = 1$$

and

$$\int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx = \sup_{\int_{B_{R_k}} (|\nabla v|^n + |v|^n) dx = 1, v \in H_0^{1,n}(B_{R_k})} \int_{B_{R_k}} \Phi(\beta_k |v|^{\frac{n}{n-1}}) dx.$$

Here, β_k is an increasing sequence tending to α_n , and R_k is an increasing sequence tending to $+\infty$.

Furthermore, u_k satisfies the following equation:

$$-div|\nabla u_k|^{n-2} \nabla u_k + u_k^{n-1} = \frac{u_k^{\frac{1}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}})}{\lambda_k},$$

where λ_k is a Lagrange multiplier.

Then, there are two possibilities. If $c_k = \max u_k$ is bounded from above, then it is easy to see that

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (\Phi(\beta_k u_k^{\frac{n}{n-1}}) - \frac{\beta_k^{n-1} u_k^n}{(n-1)!}) dx = \int_{\mathbb{R}^n} (\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1} u^n}{(n-1)!}) dx$$

where u is the weak limit of u_k . It then follows that either $\int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx$ converges to $\int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx$, or

$$S \leq \frac{\alpha_n^{n-1}}{(n-1)!}.$$

If c_k is not bounded, the key point of the proof is to show that

$$\frac{n}{n-1} \beta_k c_k^{\frac{1}{n-1}} (u_k(r_k x) - c_k) \rightarrow -n \log(1 + c_n r^{\frac{n}{n-1}}),$$

locally for a suitably chosen sequence r_k (and with $c_n = (\frac{\omega_{n-1}}{n})^{\frac{1}{n-1}}$), and that

$$c_k^{\frac{1}{n-1}} u_k \rightarrow G,$$

on any $\Omega \subset \subset \mathbb{R}^n \setminus \{0\}$, where G is some Green function. This will be done in section 3.

Then, we will get in section 4 the following

Proposition 1.3. *If S can not be attained, then*

$$S \leq \min\left\{\frac{\alpha_n^{n-1}}{(n-1)!}, \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)}\right\},$$

where $A = \lim_{r \rightarrow 0} (G(r) + \frac{1}{\alpha_n} \log r^n)$.

So, to prove the attainability, we only need to show that

$$S > \min\left\{\frac{\alpha_n^{n-1}}{(n-1)!}, \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)}\right\}.$$

In section 5, we will construct a function sequence u_ϵ such that

$$\int_{\mathbb{R}^n} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx > \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)}$$

when ϵ is sufficiently small. And in the last section we will construct, for each $n > 2$, a function sequence u_ϵ such that for ϵ sufficiently small

$$\int_{\mathbb{R}^n} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx > \frac{\alpha_n^{n-1}}{(n-1)!}.$$

Thus, together with Ruf's result of attainability in [R] for the case $n = 2$, we will get Theorem 1.2.

2 The maximizing sequence

Let $\{R_k\}$ be an increasing sequence which diverges to infinity, and $\{\beta_k\}$ an increasing sequence which converges to α_n . By compactness, we can find positive functions $u_k \in H_0^{1,n}(B_{R_k})$ with $\int_{B_{R_k}} (|\nabla u_k|^n + u_k^n) dx = 1$ such that

$$\int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx = \sup_{\int_{B_{R_k}} (|\nabla v|^n + |v|^n) dx = 1, v \in H_0^{1,n}(B_{R_k})} \int_{B_{R_k}} \Phi(\beta_k |v|^{\frac{n}{n-1}}) dx.$$

Moreover, we may assume that $\int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx = \int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx$ is increasing.

Lemma 2.1. *Let u_k as above. Then*

- a) u_k is a maximizing sequence for S ;
- b) u_k may be chosen to be radially symmetric and decreasing.

Proof. a) Let η be a cut-off function which is 1 on B_1 and 0 on $\mathbb{R}^n \setminus B_2$. Then given any $\varphi \in H^{1,n}(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} (|\nabla \varphi|^n + |\varphi|^n) dx = 1$, we have

$$\tau^n(L) := \int_{\mathbb{R}^n} (|\nabla \eta(\frac{x}{L}) \varphi|^n + |\eta(\frac{x}{L}) \varphi|^n) dx \rightarrow 1, \text{ as } L \rightarrow +\infty.$$

Hence for a fixed L and $R_k > 2L$

$$\int_{B_L} \Phi(\beta_k \left| \frac{\varphi}{\tau(L)} \right|^{\frac{n}{n-1}}) dx \leq \int_{B_{2L}} \Phi(\beta_k \left| \frac{\eta(\frac{x}{L}) \varphi}{\tau(L)} \right|^{\frac{n}{n-1}}) dx \leq \int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx$$

By the Levi Lemma, we then have

$$\int_{B_L} \Phi(\alpha_n |\frac{\varphi}{\tau(L)}|^{\frac{n}{n-1}}) dx \leq \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx.$$

Then, letting $L \rightarrow +\infty$, we get

$$\int_{\mathbb{R}^n} \Phi(\alpha_n |\varphi|^{\frac{n}{n-1}}) dx \leq \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx.$$

Hence, we get

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx = \sup_{\int_{\mathbb{R}^n} (|\nabla v|^n + |v|^n) = 1, v \in H^{1,n}(\mathbb{R}^n)} \int_{\mathbb{R}^n} \Phi(\alpha_n |v|^{\frac{n}{n-1}}) dx.$$

b) Let u_k^* be the radial rearrangement of u_k , then we have

$$\tau_k^n := \int_{B_{R_k}} (|\nabla u_k^*|^n + u_k^{*n}) dx \leq \int_{B_{R_k}} (|\nabla u_k|^n + u_k^n) dx = 1.$$

It is well-known that $\tau_k = 1$ iff u_k is radial. Since

$$\int_{B_{R_k}} \Phi(\beta_k u_k^{*\frac{n}{n-1}}) dx = \int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx,$$

we have

$$\int_{B_{R_k}} \Phi(\beta_k (\frac{u_k^*}{\tau_k})^{\frac{n}{n-1}}) dx \geq \int_{B_{R_k}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx,$$

and " $=$ " holds iff $\tau_k = 1$. Hence $\tau_k = 1$ and

$$\int_{B_{R_k}} \Phi(\beta_k u_k^{*\frac{n}{n-1}}) dx = \sup_{\int_{B_{R_k}} (|\nabla v|^n + |v|^n) = 1, v \in H_0^{1,n}(B_{R_k})} \int_{B_{R_k}} \Phi(\beta_k |v|^{\frac{n}{n-1}}) dx.$$

So, we can assume $u_k = u_k(|x|)$, and $u_k(r)$ is decreasing.

□

Assume now $u_k \rightharpoonup u$. Then, to prove Theorem 1.1 and 1.2, we only need to show that

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx = \int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx.$$

3 Blow up analysis

By the definition of u_k we have the equation

$$-div |\nabla u_k|^{n-2} \nabla u_k + u_k^{n-1} = \frac{u_k^{\frac{1}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}})}{\lambda_k}, \quad (3.1)$$

where λ_k is the constant satisfying

$$\lambda_k = \int_{B_{R_k}} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx.$$

First, we need to prove the following:

Lemma 3.1. $\inf_k \lambda_k > 0$.

Proof. Assume $\lambda_k \rightarrow 0$. Then

$$\int_{\mathbb{R}^n} u_k^n dx \leq C \int_{\mathbb{R}^n} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \leq C \lambda_k \rightarrow 0.$$

Since $u_k(|x|)$ is decreasing, we have $u_k^n(L)|B_L| \leq \int_{B_L} u_k^n \leq 1$, and then

$$u_k(L) \leq \frac{n}{\omega_n L^n}. \quad (3.2)$$

Set $\epsilon = \frac{n}{\omega_n L^n}$. Then $u_k(x) \leq \epsilon$ for any $x \notin B_L$, and hence we have, using the form of Φ , that

$$\int_{\mathbb{R}^n \setminus B_L} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx \leq C \int_{\mathbb{R}^n \setminus B_L} u_k^n dx \leq C \lambda_k \rightarrow 0.$$

And on B_L , since $u_k \rightarrow 0$ in $L^q(B_L)$ for any $q > 1$, we have by Lebesgue

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{B_L} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx &\leq \lim_{k \rightarrow +\infty} \left[\int_{B_L} C u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx + \int_{\{x \in B_L : u_k(x) \leq 1\}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx \right] \\ &\leq \lim_{k \rightarrow +\infty} C \lambda_k + \int_{B_L} \Phi(0) dx \\ &= 0. \end{aligned}$$

This is impossible. □

We denote $c_k = \max u_k = u_k(0)$. Then we have

Lemma 3.2. *If $\sup_k c_k < +\infty$, then*

- i) *Theorem 1.1 holds;*
- ii) *if S is not attained, then*

$$S \leq \frac{\alpha_n^{n-1}}{(n-1)!}.$$

Proof. If $\sup_k c_k < +\infty$, then $u_k \rightarrow u$ in $C_{loc}^1(\mathbb{R}^n)$. By (3.2), we are able to find L s.t. $u_k(x) \leq \epsilon$ for $x \notin B_L$. Then

$$\int_{\mathbb{R}^n \setminus B_L} (\Phi(\beta_k u_k^{\frac{n}{n-1}}) - \frac{\beta_k^{n-1} u_k^n}{(n-1)!}) dx \leq C \int_{\mathbb{R}^n \setminus B_L} u_k^{\frac{n^2}{n-1}} dx \leq C \epsilon^{\frac{n^2}{n-1}-n} \int_{\mathbb{R}^n} u_k^n dx \leq C \epsilon^{\frac{n^2}{n-1}-n}.$$

Letting $\epsilon \rightarrow 0$, we get

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (\Phi(\beta_k u_k^{\frac{n}{n-1}}) - \frac{\beta_k^{n-1} u_k^n}{(n-1)!}) dx = \int_{\mathbb{R}^n} (\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1} u^n}{(n-1)!}) dx.$$

Hence

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) = \int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx + \frac{\alpha_n^{n-1}}{(n-1)!} \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (u_k^n - u^n) dx. \quad (3.3)$$

When $u = 0$, we can deduce from (3.3) that

$$S \leq \frac{\alpha_n^{n-1}}{(n-1)!}.$$

Now, we assume $u \neq 0$. Set

$$\tau^n = \lim_{k \rightarrow +\infty} \frac{\int_{\mathbb{R}^n} u_k^n dx}{\int_{\mathbb{R}^n} u^n dx}.$$

By the Levi Lemma, we have $\tau \geq 1$.

Let $\tilde{u} = u(\frac{x}{\tau})$. Then, we have

$$\int_{\mathbb{R}^n} |\nabla \tilde{u}|^n dx = \int_{\mathbb{R}^n} |\nabla u|^n dx \leq \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} |\nabla u_k|^n dx,$$

and

$$\int_{\mathbb{R}^n} \tilde{u}^n dx = \tau^n \int_{\mathbb{R}^n} u^n dx = \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} u_k^n dx.$$

Then

$$\int_{\mathbb{R}^n} (|\nabla \tilde{u}|^n + \tilde{u}^n) dx \leq \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (|\nabla u_k|^n + u_k^n) dx = 1.$$

Hence, we have by (3.3)

$$\begin{aligned} S &\geq \int_{\mathbb{R}^n} \Phi(\alpha_n \tilde{u}^{\frac{n}{n-1}}) dx \\ &= \tau^n \int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx \\ &= \left[\int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx + (\tau^n - 1) \int_{\mathbb{R}^n} \frac{\alpha_n^{n-1}}{(n-1)!} u^n dx \right] + (\tau^n - 1) \int_{\mathbb{R}^n} \left(\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1}}{(n-1)!} u^n \right) dx \\ &= \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx + (\tau^n - 1) \int_{\mathbb{R}^n} \left(\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1}}{(n-1)!} u^n \right) dx \\ &= S + (\tau^n - 1) \int_{\mathbb{R}^n} \left(\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1}}{(n-1)!} u^n \right) dx \end{aligned}$$

Since $\Phi(\alpha_n u^{\frac{n}{n-1}}) - \frac{\alpha_n^{n-1}}{(n-1)!} u^n > 0$, we have $\tau = 1$, and then

$$S = \int_{\mathbb{R}^n} \Phi(\alpha_n u^{\frac{n}{n-1}}) dx.$$

So, u is an extremal function. □

From now on, we assume $c_k \rightarrow +\infty$. We perform a blow-up procedure:
We define

$$r_k^n = \frac{\lambda_k}{c_k^{\frac{n}{n-1}} e^{\beta_k c_k^{\frac{n}{n-1}}}}.$$

By (3.2) we can find a sufficiently large L such that $u_k \leq 1$ on $\mathbb{R}^n \setminus B_L$. Then

$$\int_{B_L} |\nabla(u_k - u_k(L))^+|^n dx \leq 1$$

and hence, by (1.1), we have

$$\int_{B_L} e^{\alpha_n [(u_k - u_k(L))^+]^{\frac{n}{n-1}}} \leq C(L).$$

Clearly, for any $p < \alpha_n$ we can find a constant $C(p)$, s.t.

$$p u_k^{\frac{n}{n-1}} \leq \alpha_n [(u_k - u_k(L))^+]^{\frac{n}{n-1}} + C(p),$$

and then we get

$$\int_{B_L} e^{p u_k^{\frac{n}{n-1}}} dx < C = C(L, p).$$

Hence,

$$\begin{aligned} \lambda_k e^{-\frac{\beta_k}{2} c_k^{\frac{n}{n-1}}} &= e^{-\frac{\beta_k}{2} c_k^{\frac{n}{n-1}}} \left[\int_{\mathbb{R}^n \setminus B_L} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx + \int_{B_L} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \right] \\ &\leq C \int_{\mathbb{R}^n \setminus B_L} u_k^n dx e^{-\frac{\beta_k}{2} c_k^{\frac{n}{n-1}}} + \int_{B_L} e^{\frac{\beta_k}{2} u_k^{\frac{n}{n-1}}} u_k^{\frac{n}{n-1}} dx. \end{aligned}$$

Since u_k converges strongly in $L^q(B_L)$ for any $q > 1$, we get $\lambda_k \leq C e^{\frac{\beta_k}{2} c_k^{\frac{n}{n-1}}}$, and hence

$$r_k^n \leq C e^{-\frac{\beta_k}{2} c_k^{\frac{n}{n-1}}}.$$

Now, we set

$$v_k(x) = u_k(r_k x), \quad w_k(x) = \frac{n}{n-1} \beta_k c_k^{\frac{1}{n-1}} (v_k - c_k),$$

where v_k and w_k are defined on $\Omega_k = \{x \in \mathbb{R}^n : r_k x \in B_1\}$. Using the definition of r_k^n and (3.1) we have

$$-div |\nabla w_k|^{n-2} \nabla w_k = \frac{v_k^{\frac{1}{n-1}}}{c_k^{\frac{1}{n-1}}} \left(\frac{n}{n-1} \beta_k \right)^{n-1} e^{\beta_k (v_k^{\frac{n}{n-1}} - c_k^{\frac{n}{n-1}})} + O(r_k^n c_k^n).$$

By Theorem 7 in [S], we know that $osc_{B_R} \omega_k \leq C(R)$ for any $R > 0$. Then from the result in [T] (or [D]), it follows that $\|w_k\|_{C^{1,\delta}(B_R)} < C(R)$. Therefore w_k converges in C_{loc}^1 and $v_k - c_k \rightarrow 0$ in C_{loc}^1 .

Since

$$v_k^{\frac{n}{n-1}} = c_k^{\frac{n}{n-1}} \left(1 + \frac{v_k - c_k}{c_k} \right)^{\frac{n}{n-1}} = c_k^{\frac{n}{n-1}} \left(1 + \frac{n}{n-1} \frac{v_k - c_k}{c_k} + O\left(\frac{1}{c_k^2}\right) \right),$$

we get $\beta_k (v_k^{\frac{n}{n-1}} - c_k^{\frac{n}{n-1}}) \rightarrow w$ in C_{loc}^0 , and so we have

$$-div |\nabla w|^{n-2} \nabla w = \left(\frac{n \alpha_n}{n-1} \right)^{n-1} e^w, \quad (3.4)$$

with

$$w(0) = 0 = \max w.$$

Since ω is radially symmetric and decreasing, it is easy to see that (3.4) has only one solution. We can check that

$$w(x) = -n \log(1 + c_n |x|^{\frac{n}{n-1}}), \quad \text{and} \quad \int_{\mathbb{R}^n} e^w dx = 1,$$

where $c_n = (\frac{\omega_{n-1}}{n})^{\frac{1}{n-1}}$. Then,

$$\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_{Lr_k}} \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} e^{\beta_k u_k^{\frac{n}{n-1}}} dx = \lim_{L \rightarrow +\infty} \int_{B_L} e^w dx = 1. \quad (3.5)$$

For $A > 1$, let $u_k^A = \min\{u_k, \frac{c_k}{A}\}$. We have

Lemma 3.3. *For any $A > 1$, there holds*

$$\limsup_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (|\nabla u_k^A|^n + |u_k^A|^n) dx \leq \frac{1}{A}. \quad (3.6)$$

Proof. Since $|\{x : u_k \geq \frac{c_k}{A}\}| |\frac{c_k}{A}|^n \leq \int_{\{u_k \geq \frac{c_k}{A}\}} u_k^n \leq 1$, we can find a sequence $\rho_k \rightarrow 0$ s.t.

$$\{x : u_k \geq \frac{c_k}{A}\} \subset B_{\rho_k}.$$

Since u_k converges in $L^p(B_1)$ for any $p > 1$, we have

$$\lim_{k \rightarrow +\infty} \int_{\{u_k > \frac{c_k}{A}\}} |u_k^A|^p dx \leq \lim_{k \rightarrow +\infty} \int_{\{u_k > \frac{c_k}{A}\}} u_k^p dx = 0,$$

and

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (u_k - \frac{c_k}{A})^+ u_k^p dx = 0$$

for any $p > 0$.

Hence, testing equation (3.1) with $(u_k - \frac{c_k}{A})^+$, we have

$$\begin{aligned} \int_{\mathbb{R}^n} \left(|\nabla (u_k - \frac{c_k}{A})^+|^n + (u_k - \frac{c_k}{A})^+ u_k^{n-1} \right) dx &= \int_{\mathbb{R}^n} (u_k - \frac{c_k}{A})^+ \frac{u_k^{\frac{1}{n-1}}}{\lambda_k} e^{\beta_k u_k^{\frac{n}{n-1}}} dx + o(1) \\ &\geq \int_{B_{Lr_k}} (u_k - \frac{c_k}{A})^+ \frac{u_k^{\frac{1}{n-1}}}{\lambda_k} e^{\beta_k u_k^{\frac{n}{n-1}}} dx + o(1) \\ &= \int_{B_L} \frac{v_k - c_k/A}{c_k} \left(\frac{v_k - c_k}{c_k} + 1 \right)^{\frac{1}{n-1}} e^{w_k + o(1)} dx + o(1). \end{aligned}$$

Hence

$$\liminf_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \left(|\nabla (u_k - \frac{c_k}{A})^+|^n + (u_k - \frac{c_k}{A})^+ u_k^{n-1} \right) dx \geq \frac{A-1}{A} \int_{B_L} e^w dx.$$

Letting $L \rightarrow +\infty$, we get

$$\liminf_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \left(|\nabla(u_k - \frac{c_k}{A})^+|^n + (u_k - \frac{c_k}{A})^+ u_k^{n-1} \right) dx \geq \frac{A-1}{A}.$$

Now observe that

$$\begin{aligned} \int_{\mathbb{R}^n} (|\nabla u_k^A|^n + |u_k^A|^n) dx &= 1 - \int_{\mathbb{R}^n} \left(|\nabla(u_k - \frac{c_k}{A})^+|^n + (u_k - \frac{c_k}{A})^+ u_k^{n-1} \right) dx \\ &\quad + \int_{\mathbb{R}^n} (u_k - \frac{c_k}{A})^+ u_k^{n-1} dx - \int_{\{u_k > \frac{c_k}{A}\}} u_k^n dx + \int_{\{u_k > \frac{c_k}{A}\}} |u_k^A|^n dx \\ &\leq 1 - (1 - \frac{1}{A}) + o(1). \end{aligned}$$

Hence, we get this Lemma. □

Corollary 3.4. *We have*

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n \setminus B_\delta} (|\nabla u_k|^n + u_k^n) dx = 0 ,$$

for any $\delta > 0$, and then $u = 0$.

Proof. Letting $A \rightarrow +\infty$, then for any constant c , we have

$$\int_{\{u_k \leq c\}} (|\nabla u_k|^n + u_k^n) dx \rightarrow 0.$$

So we get this Corollary. □

Lemma 3.5. *We have*

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx \leq \lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_{Lr_k}} (e^{\beta_k u_k^{\frac{n}{n-1}}} - 1) dx = \limsup_{k \rightarrow \infty} \frac{\lambda_k}{c_k^{\frac{n}{n-1}}}, \quad (3.7)$$

and consequently

$$\frac{\lambda_k}{c_k} \rightarrow +\infty , \quad \text{and} \quad \sup_k \frac{c_k^{\frac{n}{n-1}}}{\lambda_k} < +\infty . \quad (3.8)$$

Proof. We have

$$\begin{aligned} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx &\leq \int_{\{u_k \leq \frac{c_k}{A}\}} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx + \int_{\{u_k > \frac{c_k}{A}\}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \\ &\leq \int_{\mathbb{R}^n} \Phi(\beta_k (u_k^A)^{\frac{n}{n-1}}) dx + A^{\frac{n}{n-1}} \frac{\lambda_k}{c_k^{\frac{n}{n-1}}} \int_{\mathbb{R}^n} \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx . \end{aligned}$$

Applying (3.2), we can find L such that $u_k \leq 1$ on $\mathbb{R}^n \setminus B_L$. Then by Corollary 3.4 and the form of Φ , we have

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n \setminus B_L} \Phi(p\beta_k(u_k^A)^{\frac{n}{n-1}}) dx \leq \lim_{k \rightarrow \infty} C(p) \int_{\mathbb{R}^n \setminus B_L} u_k^n dx = 0 \quad (3.9)$$

for any $p > 0$.

Since by Lemma 3.3 $\limsup_{k \rightarrow +\infty} \int_{\mathbb{R}^n} (|\nabla u_k^A|^n + |u_k^A|^n) dx \leq \frac{1}{A} < 1$ when $A > 1$, it follows from (1.1) that

$$\sup_k \int_{B_L} e^{p' \beta_k((u_k^A - u_k(L))^+)^{\frac{n}{n-1}}} dx < +\infty$$

for any $p' < A^{\frac{1}{n-1}}$. Since for any $p < p'$

$$p(u_k^A)^{\frac{n}{n-1}} \leq p'((u_k^A - u_k(L))^+)^{\frac{n}{n-1}} + C(p, p'),$$

we have

$$\sup_k \int_{B_L} \Phi(p\beta_k(u_k^A)^{\frac{n}{n-1}}) dx < +\infty \quad (3.10)$$

for any $p < A^{\frac{1}{n-1}}$. Then on B_L , by the weak compactness of Banach space, we get

$$\lim_{k \rightarrow +\infty} \int_{B_L} \Phi(\beta_k(u_k^A)^{\frac{n}{n-1}}) dx = \int_{B_L} \Phi(0) dx = 0.$$

Hence we have

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx &\leq \lim_{L \rightarrow \infty} \lim_{k \rightarrow +\infty} A^{\frac{n}{n-1}} \frac{\lambda_k}{c_k^{\frac{n}{n-1}}} \int_{B_L} \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx + C\epsilon \\ &= \lim_{k \rightarrow +\infty} A^{\frac{n}{n-1}} \frac{\lambda_k}{c_k^{\frac{n}{n-1}}} + C\epsilon. \end{aligned}$$

As $A \rightarrow 1$ and $\epsilon \rightarrow 0$ we obtain (3.7).

If $\frac{\lambda_k}{c_k}$ was bounded or $\sup_k \frac{c_k^{\frac{n}{n-1}}}{\lambda_k} = +\infty$, it would follow from (3.7) that

$$\sup_{\int_{\mathbb{R}^n} (|\nabla v|^n + |v|^n) dx = 1, v \in H^{1,n}(\mathbb{R}^n)} \int_{\mathbb{R}^n} \Phi(\alpha_n |v|^{\frac{n}{n-1}}) dx = 0,$$

which is impossible. □

Lemma 3.6. *We have that $c_k \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}})$ converges to δ_0 weakly, i.e. for any $\varphi \in \mathcal{D}(\mathbb{R}^n)$ we have*

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \varphi c_k \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx = \varphi(0).$$

Proof. Suppose $\text{supp } \varphi \subset B_\rho$. We split the integral

$$\begin{aligned} \int_{B_\rho} \varphi \frac{c_k u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx &\leq \int_{\{u_k \geq \frac{c_k}{A}\} \setminus B_{Lr_k}} \cdots + \int_{B_{Lr_k}} \cdots + \int_{\{u_k < \frac{c_k}{A}\}} \cdots \\ &= I_1 + I_2 + I_3 . \end{aligned}$$

We have

$$I_1 \leq A \|\varphi\|_{C^0} \int_{\mathbb{R}^n \setminus B_{Lr_k}} \frac{u_k^{\frac{n}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx = A \|\varphi\|_{C^0} (1 - \int_{B_L} e^{w_k + o(1)} dx) ,$$

and

$$I_2 = \int_{B_L} \varphi(r_k x) \frac{c_k (c_k + (v_k - c_k))^{\frac{1}{n-1}}}{c_k^{\frac{n}{n-1}}} e^{w_k + o(1)} dx = \varphi(0) \int_{B_L} e^w dx + o(1) = \varphi(0) + o(1) .$$

By (3.9) and (3.10) we have

$$\int_{\mathbb{R}^n} \Phi(p\beta_k |u_k^A|^{\frac{n}{n-1}}) dx < C$$

for any $p < A^{\frac{1}{n-1}}$. We set $\frac{1}{q} + \frac{1}{p} = 1$. Then we get by (3.8)

$$I_3 = \int_{\{u_k \leq \frac{c_k}{A}\}} \varphi c_k \frac{u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \leq \frac{c_k}{\lambda_k} \|\varphi\|_{C^0} \|u_k^{\frac{1}{n-1}}\|_{L^q(\mathbb{R}^n)} \|e^{\beta_k |u_k^A|^{\frac{n}{n-1}}}\|_{L^p(\mathbb{R}^n)} \rightarrow 0.$$

Letting $L \rightarrow +\infty$, we deduce now that

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n} \varphi \frac{c_k u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx = \varphi(0).$$

□

Proposition 3.7. *On any $\Omega \subset \subset \mathbb{R}^n \setminus \{0\}$, we have that $c_k^{\frac{1}{n-1}} u_k$ converges to G in $C^1(\Omega)$, where $G \in C_{loc}^{1,\alpha}(\mathbb{R}^n \setminus \{0\})$ satisfies the following equation:*

$$-div|\nabla G|^{n-2} \nabla G + G^{n-1} = \delta_0 . \quad (3.11)$$

Proof. We set $U_k = c_k^{\frac{1}{n-1}} u_k$, which satisfy by (3.1) the equations:

$$-div|\nabla U_k|^{n-2} \nabla U_k + U_k^{n-1} = \frac{c_k u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) . \quad (3.12)$$

For our purpose, we need to prove that

$$\int_{B_R} |U_k|^q dx \leq C(q, R) ,$$

where $C(q, R)$ does not depend on k . We use the idea in [St] to prove this statement.

Set $\Omega_t = \{0 \leq U_k \leq t\}$, $U_k^t = \min\{U_k, t\}$. Then we have

$$\int_{\Omega_t} (|\nabla U_k^t|^n + |U_k^t|^n) dx \leq \int_{\mathbb{R}^n} (-U_k^t \Delta_n U_k + U_k^t U_k^{n-1}) = \int_{\mathbb{R}^n} U_k^t \frac{c_k u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \leq 2t.$$

Let η be a radially symmetric cut-off function which is 1 on B_R and 0 on B_{2R}^c . Then,

$$\int_{B_{2R}} |\nabla \eta U_k^t|^n dx \leq C_1(R) + C_2(R)t.$$

Then, when t is bigger than $\frac{C_1(R)}{C_2(R)}$, we have

$$\int_{B_{2R}} |\nabla \eta U_k^t|^n dx \leq 2C_2(R)t.$$

Set ρ such that $U_k(\rho) = t$. Then we have

$$\inf \left\{ \int_{B_{2R}} |\nabla v|^n dx : v \in H_0^{1,n}(B_{2R}) \text{ and } v|_{B_\rho} = t \right\} \leq 2C_2(R)t.$$

On the other hand, the inf is achieved by $-t \log \frac{|x|}{2R} / \log \frac{2R}{\rho}$. By a direct computation, we have

$$\frac{\omega_{n-1} t^{n-1}}{(\log \frac{2R}{\rho})^{n-1}} \leq 2C_2(R),$$

and hence for any $t > \frac{C_1(R)}{C_2(R)}$

$$|\{x \in B_{2R} : U_k \geq t\}| = |B_\rho| \leq C_3(R) e^{-A(R)t},$$

where $A(R)$ is a constant only depending on R . Then, for any $\delta < A$,

$$\int_{B_R} e^{\delta U_k} dx \leq \sum_{m=0}^{\infty} \mu(\{m \leq U_k \leq m+1\}) e^{\delta(m+1)} \leq \sum_{m=0}^{\infty} e^{-(A-\delta)m} e^{\delta} \leq C.$$

Then, testing the equation (3.12) with the function $\log \frac{1+2(U_k - U_k(R))^+}{1+(U_k - U_k(R))^+}$, we get

$$\begin{aligned} & \int_{B_R} \frac{|\nabla U_k|^n}{(1+U_k - U_k(R))(1+2U_k - 2U_k(R))} dx \\ & \leq \log 2 \int_{B_R} \frac{c_k u_k^{\frac{1}{n-1}}}{\lambda_k} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx - \int_{B_R} U_k^{n-1} \log \frac{1+2(U_k - U_k(R))}{1+(U_k - U_k(R))} dx \leq C. \end{aligned}$$

Given $q < n$, by Young's Inequality, we have

$$\begin{aligned} \int_{B_R} |\nabla U_k|^q dx & \leq \int_{B_R} \left[\frac{|\nabla U_k|^n}{(1+U_k - U_k(R))(1+2U_k - 2U_k(R))} + ((1+U_k)(1+2U_k))^{\frac{n}{n-q}} \right] dx \\ & \leq \int_{B_R} \left[\frac{|\nabla U_k|^n}{(1+U_k - U_k(R))(1+2U_k - 2U_k(R))} + C e^{\delta U_k} \right] dx. \end{aligned}$$

Hence, we are able to assume that U_k converges to a function G weakly in $H^{1,p}(B_R)$ for any R and $p < n$. Applying Lemma 3.6, we get (3.11).

Hence U_k is bounded in $L^q(\Omega)$ for any $q > 0$. By Corollary 3.4 and Theorem A, $e^{\beta_k u_k^{\frac{n}{n-1}}}$ is also bounded in $L^q(\Omega)$ for any $q > 0$. Then, applying Theorem 2.8 in [S], and the main result in [T] (or [D]), we get $\|U_k\|_{C^{1,\alpha}(\Omega)} \leq C$. So, U_k converges to G in $C^1(\Omega)$. $\square$

For the Green function G we have the following results:

Lemma 3.8. $G \in C_{loc}^{1,\alpha}(\mathbb{R}^n \setminus \{0\})$ and near 0 we can write

$$G = -\frac{1}{\alpha_n} \log r^n + A + O(r^n \log^n r); \quad (3.13)$$

here, A is a constant. Moreover, for any $\delta > 0$, we have

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n \setminus B_\delta} (|\nabla c_k^{\frac{1}{n-1}} u_k|^n + (c_k^{\frac{1}{n-1}} u_k)^n) dx &= \int_{\mathbb{R}^n \setminus B_\delta} (|\nabla G|^n + |G|^n) dx \\ &= G(\delta) \left(1 - \int_{B_\delta} G^{n-1} dx\right). \end{aligned}$$

Proof. Slightly modifying the proof in [K-L], we can prove

$$G = -\frac{1}{\alpha_n} \log r^n + A + o(1).$$

One can refer to [L2] for details. Further, testing the equation (3.12) with 1, we get

$$\omega_{n-1} G'(r)^{n-1} r^{n-1} = \int_{\partial B_r} |\nabla G|^{n-2} \frac{\partial G}{\partial n} = 1 - \int_{B_r} G^{n-1} dx = 1 + O(r^n \log^{n-1} r).$$

Then, we get (3.13).

We have

$$\int_{\mathbb{R}^n \setminus B_\delta} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx \leq C \int_{\mathbb{R}^n \setminus B_\delta} u_k^n dx \rightarrow 0. \quad (3.14)$$

Recall that $U_k \in H_0^{1,n}(B_{R_k})$. By equation (3.12) we get

$$\int_{\mathbb{R}^n \setminus B_\delta} (|\nabla U_k|^n + U_k^n) dx = \frac{c_k^{\frac{n}{n-1}}}{\lambda_k} \int_{\mathbb{R}^n \setminus B_\delta} u_k^{\frac{n}{n-1}} \Phi'(\beta_k u_k^{\frac{n}{n-1}}) dx - \int_{\partial B_\delta} \frac{\partial U_k}{\partial n} |\nabla U_k|^{n-2} U_k dS.$$

By (3.14) and (3.8) we then get

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^n \setminus B_\delta} (|\nabla U_k|^n + U_k^n) dx &= - \lim_{k \rightarrow +\infty} \int_{\partial B_\delta} \frac{\partial U_k}{\partial n} |\nabla U_k|^{n-2} U_k dS \\ &= -G(\delta) \int_{\partial B_\delta} \frac{\partial G}{\partial n} |\nabla G|^{n-2} dS \\ &= G(\delta) \left(1 - \int_{B_\delta} G^{n-1} dx\right). \end{aligned}$$

□

We are now in the position to complete the proof of Theorem 1.1: We have seen in (3.9) that

$$\int_{\mathbb{R}^n \setminus B_R} \Phi(\beta_k u_k^{\frac{n}{n-1}}) dx \leq C.$$

So, we only need to prove on B_R ,

$$\int_{B_R} e^{\beta_k u_k^{\frac{n}{n-1}}} dx < C$$

The classical Trudinger-Moser inequality implies that

$$\int_{B_R} e^{\beta_k ((u_k - u_k(R))^+)^{\frac{n}{n-1}}} dx < C = C(R).$$

By Proposition 3.7, $u_k(R) = O(\frac{1}{c_k^{\frac{n-1}{n}}})$, and hence we have

$$u_k^{\frac{n}{n-1}} \leq ((u_k - u_k(R))^+ + u_k(R))^{\frac{n}{n-1}} \leq ((u_k - u_k(R))^+)^{\frac{n}{n-1}} + C_1,$$

Then, we get

$$\int_{B_R} e^{\beta_k u_k^{\frac{n}{n-1}}} \leq C'.$$

□

4 The proof of Proposition 1.3

We will use a result of Carleson and Chang (see [C-C]):

Lemma 4.1. *Let B be the unit ball in $\mathbb{R}^n$. Assume that u_k is a sequence in $H_0^{1,n}(B)$ with $\int_B |\nabla u_k|^n dx = 1$. If $u_k \rightarrow 0$, then*

$$\limsup_{k \rightarrow +\infty} \int_B (e^{\alpha_n |u_k|^{\frac{n}{n-1}}} - 1) dx \leq |B| e^{1+1/2+\dots+1/(n-1)}.$$

Proof of Proposition 1.3: Set $u'_k(x) = \frac{(u_k(x) - u_k(\delta))^+}{\|\nabla u_k\|_{L^n(B_\delta)}}$ which is in $H_0^{1,n}(B_\delta)$. Then by the result of Carleson and Chang, we have

$$\limsup_{k \rightarrow +\infty} \int_{B_\delta} e^{\beta_k u'_k{}^{\frac{n}{n-1}}} \leq |B_\delta| (1 + e^{1+1/2+\dots+1/(n-1)}).$$

By Lemma 3.8, we have

$$\int_{\mathbb{R}^n \setminus B_\delta} (|\nabla c_k^{\frac{1}{n-1}} u_k|^n + (c_k^{\frac{1}{n-1}} u_k)^n) dx \rightarrow G(\delta) (1 - \int_{B_\delta} G^{n-1} dx),$$

and therefore we get

$$\int_{B_\delta} |\nabla u_k|^n dx = 1 - \int_{\mathbb{R}^n \setminus B_\delta} (|\nabla u_k|^n + u_k^n) dx - \int_{B_\delta} u_k^n dx = 1 - \frac{G(\delta) + \epsilon_k(\delta)}{c_k^{\frac{n}{n-1}}}, \quad (4.1)$$

where $\lim_{\delta \rightarrow 0} \lim_{k \rightarrow +\infty} \epsilon_k(\delta) = 0$.

By (3.9) in Lemma 3.5 we have

$$\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_\rho \setminus B_{Lr_k}} e^{\beta_k u_k^{\frac{n}{n-1}}} dx = |B_\rho|,$$

for any $\rho < \delta$. Furthermore, on B_ρ we have by (4.1)

$$\begin{aligned} (u'_k)^{\frac{n}{n-1}} &\leq \frac{u_k^{\frac{n}{n-1}}}{\left(1 - \frac{G(\delta) + \epsilon_k(\delta)}{c_k^{\frac{n}{n-1}}}\right)^{\frac{1}{n-1}}} = u_k^{\frac{n}{n-1}} \left(1 + \frac{1}{n-1} \frac{G(\delta) + \epsilon_k(\delta)}{c_k^{\frac{n}{n-1}}} + O\left(\frac{1}{c_k^{\frac{2n}{n-1}}}\right)\right) \\ &= u_k^{\frac{n}{n-1}} + \frac{1}{n-1} G(\delta) \left(\frac{u_k}{c_k}\right)^{\frac{n}{n-1}} + O(c_k^{\frac{-n}{n-1}}) \\ &\leq u_k^{\frac{n}{n-1}} - \frac{\log \delta^n}{(n-1)\alpha_n}. \end{aligned}$$

Then we have

$$\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_\rho \setminus B_{Lr_k}} e^{\beta_k u'_k^{\frac{n}{n-1}}} dx \leq O(\delta^{-n}) \lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_\rho \setminus B_{Lr_k}} e^{\beta_k u_k^{\frac{n}{n-1}}} dx \rightarrow |B_\rho| O(\delta^{-n}).$$

Since $u'_k \rightarrow 0$ on $B_\delta \setminus B_\rho$, we get

$$\lim_{k \rightarrow +\infty} \int_{B_\delta \setminus B_\rho} (e^{\beta_k u'_k^{\frac{n}{n-1}}} - 1) dx = 0,$$

then

$$0 \leq \lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_\delta \setminus B_{Lr_k}} (e^{\beta_k u'_k^{\frac{n}{n-1}}} - 1) dx \leq |B_\rho| O(\delta^{-n}).$$

Letting $\rho \rightarrow 0$, we get

$$\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_\delta \setminus B_{Lr_k}} (e^{\beta_k u'_k^{\frac{n}{n-1}}} - 1) dx = 0.$$

So, we have

$$\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_{Lr_k}} (e^{\beta_k u'_k^{\frac{n}{n-1}}} - 1) dx \leq e^{1+1/2+\dots+1/(n-1)} |B_\delta|.$$

Now, we fix an L . Then for any $x \in B_{Lr_k}$, we have

$$\begin{aligned}
\beta_k u_k^{\frac{n}{n-1}} &= \beta_k \left(\frac{u_k}{\|\nabla u_k\|_{L^n(B_\delta)}} \right)^{\frac{n}{n-1}} \left(\int_{B_\delta} |\nabla u_k|^n dx \right)^{\frac{1}{n-1}} \\
&= \beta_k \left(u'_k + \frac{u_k(\delta)}{\|\nabla u_k\|_{L^n(B_\delta)}} \right)^{\frac{n}{n-1}} \left(\int_{B_\delta} |\nabla u_k|^n dx \right)^{\frac{1}{n-1}} \\
&\quad \left(\text{using that } u_k(\delta) = O\left(\frac{1}{c_k^{\frac{1}{n-1}}}\right) \text{ and } \|\nabla u_k\|_{L^n(B_\delta)} = 1 + O\left(\frac{1}{c_k^{\frac{1}{n-1}}}\right) \right) \\
&= \beta_k \left(u'_k + u_k(\delta) + O\left(\frac{1}{c_k^{\frac{n+1}{n-1}}}\right) \right)^{\frac{n}{n-1}} \left(\int_{B_\delta} |\nabla u_k|^n dx \right)^{\frac{1}{n-1}} \\
&= \beta_k u'_k^{\frac{n}{n-1}} \left(1 + \frac{u_k(\delta)}{u'_k} + O\left(\frac{1}{c_k^{\frac{2n}{n-1}}}\right) \right)^{\frac{n}{n-1}} \left(1 - \frac{G(\delta) + \epsilon_k(\delta)}{c_k^{\frac{n}{n-1}}} \right)^{\frac{1}{n-1}} \\
&= \beta_k u'_k^{\frac{n}{n-1}} \left[1 + \frac{n}{n-1} \frac{u_k(\delta)}{u'_k} - \frac{1}{n-1} \frac{G(\delta) + \epsilon_k(\delta)}{c_k^{\frac{n}{n-1}}} + O\left(\frac{1}{c_k^{\frac{2n}{n-1}}}\right) \right].
\end{aligned}$$

It is easy to check that

$$\frac{u'_k(r_k x)}{c_k} \rightarrow 1, \quad \text{and} \quad (u'_k(r_k x))^{\frac{1}{n-1}} u_k(\delta) \rightarrow G(\delta).$$

So, we get

$$\begin{aligned}
\lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} \int_{B_{Lr_k}} (e^{\beta_k u_k^{\frac{n}{n-1}}} - 1) dx &= \lim_{L \rightarrow +\infty} \lim_{k \rightarrow +\infty} e^{\alpha_n G(\delta)} \int_{B_{Lr_k}} (e^{\beta_k u'_k \frac{n}{n-1}} - 1) dx \\
&\leq e^{\alpha_n G(\delta)} \delta^n \frac{\omega_{n-1}}{n} e^{1+1/2+\dots+1/(n-1)} \\
&= e^{\alpha_n (-\frac{1}{\alpha_n} \log \delta^n + A + O(\delta^n \log^n \delta))} \delta^n \frac{\omega_{n-1}}{n} e^{1+1/2+\dots+1/(n-1)}.
\end{aligned}$$

Letting $\delta \rightarrow 0$, then the above inequality together with Lemma 3.2 imply Proposition 1.3.

5 The test function 1

In this section, we will construct a function sequence $\{u_\epsilon\} \subset H^{1,n}(\mathbb{R}^n)$ with $\|u_\epsilon\|_{H^{1,n}} = 1$ which satisfies

$$\int_{\mathbb{R}^n} \Phi(\alpha_n |u_\epsilon|^{\frac{n}{n-1}}) dx > \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)},$$

for $\epsilon > 0$ sufficiently small.

Let

$$u_\epsilon = \begin{cases} C - \frac{(n-1) \log(1 + c_n |\frac{x}{\epsilon}|^{\frac{n}{n-1}}) + \Lambda_\epsilon}{\alpha_n C^{\frac{1}{n-1}}} & |x| \leq L\epsilon \\ \frac{G(|x|)}{C^{\frac{1}{n-1}}} & |x| > L\epsilon, \end{cases}$$

where Λ_ϵ, C and L are functions of ϵ (which will be defined later, by (5.1), (5.2), (5.5)) which satisfy

- i) $L \rightarrow +\infty$, $C \rightarrow +\infty$, and $L\epsilon \rightarrow 0$, as $\epsilon \rightarrow 0$;
- ii) $C - \frac{(n-1) \log(1+c_n L^{\frac{n}{n-1}}) + \Lambda_\epsilon}{\alpha_n C^{\frac{1}{n-1}}} = \frac{G(L\epsilon)}{C^{\frac{1}{n-1}}}$;
- iii) $\frac{\log L}{C^{\frac{n}{n-1}}} \rightarrow 0$, as $\epsilon \rightarrow 0$.

We use the normalization of u_ϵ to obtain information on Λ_ϵ, C and L . We have

$$\begin{aligned}
\int_{\mathbb{R}^n \setminus B_{L\epsilon}} (|\nabla u_\epsilon|^n + u_\epsilon^n) dx &= \frac{1}{C^{\frac{n}{n-1}}} \left(\int_{B_{L\epsilon}^c} |\nabla G|^n dx + \int_{B_{L\epsilon}^c} G^n dx \right) \\
&= \frac{1}{C^{\frac{n}{n-1}}} \int_{\partial B_{L\epsilon}} G(L\epsilon) |\nabla G|^{n-2} \frac{\partial G}{\partial n} dS \\
&= \frac{G(L\epsilon) - G(L\epsilon) \int_{B_{L\epsilon}} G^{n-1} dx}{C^{\frac{n}{n-1}}} .
\end{aligned}$$

and

$$\begin{aligned}
\int_{B_{L\epsilon}} |\nabla u_\epsilon|^n dx &= \frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} \int_0^{c_n L^{\frac{n}{n-1}}} \frac{u^{n-1}}{(1+u)^n} du \\
&= \frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} \int_0^{c_n L^{\frac{n}{n-1}}} \frac{((1+u)-1)^{n-1}}{(1+u)^n} du \\
&= \frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} \sum_{k=0}^{n-2} \frac{C_{n-1}^k (-1)^{n-1-k}}{n-k-1} \\
&\quad + \frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} \log(1 + c_n L^{\frac{n}{n-1}}) + O\left(\frac{1}{L^{\frac{n}{n-1}} C^{\frac{n}{n-1}}}\right) \\
&= -\frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} (1 + 1/2 + 1/3 + \dots + 1/(n-1)) \\
&\quad + \frac{n-1}{\alpha_n C^{\frac{n}{n-1}}} \log(1 + c_n L^{\frac{n}{n-1}}) + O\left(\frac{1}{L^{\frac{n}{n-1}} C^{\frac{n}{n-1}}}\right) ,
\end{aligned}$$

where we used the fact

$$-\sum_{k=0}^{n-2} \frac{C_{n-1}^k (-1)^{n-1-k}}{n-k-1} = 1 + \frac{1}{2} + \dots + \frac{1}{n-1} .$$

It is easy to check that

$$\int_{B_{L\epsilon}} |u_\epsilon|^n dx = O((L\epsilon)^n C^n \log L) ,$$

and thus we get

$$\begin{aligned}
\int_{\mathbb{R}^n} (|\nabla u_\epsilon|^n + u_\epsilon^n) dx &= \frac{1}{\alpha_n C^{\frac{n}{n-1}}} \left\{ -(n-1) \left(1 + 1/2 + \dots + 1/(n-1) \right) + \alpha_n A \right. \\
&\quad \left. + (n-1) \log(1 + c_n L^{\frac{n}{n-1}}) - \log(L\epsilon)^n + \phi \right\} ,
\end{aligned}$$

where

$$\phi = O\left((L\epsilon)^n C^n \log L + (L\epsilon)^n \log^n L\epsilon + L^{\frac{-n}{n-1}}\right).$$

Setting $\int_{\mathbb{R}^n} (|\nabla u_\epsilon|^n + u_\epsilon^n) dx = 1$, we obtain

$$\begin{aligned} \alpha_n C^{\frac{n}{n-1}} &= -(n-1)\left(1 + 1/2 + \cdots + 1/(n-1)\right) + \alpha_n A + \log \frac{(1+c_n L^{\frac{n}{n-1}})^{n-1}}{L^n} - \log \epsilon^n + \phi \\ &= -(n-1)\left(1 + 1/2 + \cdots + 1/(n-1)\right) + \alpha_n A + \log \frac{\omega_{n-1}}{n} - \log \epsilon^n + \phi. \end{aligned} \quad (5.1)$$

By ii) we have

$$\alpha_n C^{\frac{n}{n-1}} - (n-1) \log(1 + c_n L^{\frac{n}{n-1}}) + \Lambda_\epsilon = \alpha G(L\epsilon)$$

and hence

$$-(n-1)\left(1 + 1/2 + \cdots + 1/(n-1)\right) + \alpha_n A - \log(L\epsilon)^n + \phi + \Lambda_\epsilon = \alpha G(L\epsilon);$$

this implies that

$$\Lambda_\epsilon = -(n-1)\left(1 + 1/2 + \cdots + 1/(n-1)\right) + \phi. \quad (5.2)$$

Next, we compute $\int_{B_{L\epsilon}} e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} dx$.

Clearly, $\varphi(t) = |1-t|^{\frac{n}{n-1}} + \frac{n}{n-1}t$ is increasing when $0 \leq t \leq 1$ and decreasing when $t \leq 0$, then

$$|1-t|^{\frac{n}{n-1}} \geq 1 - \frac{n}{n-1}t, \quad \text{when } |t| < 1.$$

Thus we have by ii), for any $x \in B_{L\epsilon}$

$$\begin{aligned} \alpha_n u_\epsilon^{\frac{n}{n-1}} &= \alpha_n C^{\frac{n}{n-1}} \left| 1 - \frac{(n-1) \log(1 + c_n |\frac{x}{\epsilon}|^{\frac{n}{n-1}}) + \Lambda_\epsilon}{\alpha_n C^{\frac{n}{n-1}}} \right|^{\frac{n}{n-1}} \\ &\geq \alpha_n C^{\frac{n}{n-1}} \left(1 - \frac{n}{n-1} \frac{(n-1) \log(1 + c_n |\frac{x}{\epsilon}|^{\frac{n}{n-1}}) + \Lambda_\epsilon}{\alpha_n C^{\frac{n}{n-1}}} \right). \end{aligned} \quad (5.3)$$

Then we have

$$\begin{aligned} \int_{B_{L\epsilon}} e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} dx &\geq \int_{B_{L\epsilon}} e^{\alpha_n C^{\frac{n}{n-1}} - n \log(1 + c_n |\frac{x}{\epsilon}|^{\frac{n}{n-1}}) - \frac{n}{n-1} \Lambda_\epsilon} \\ &= e^{\alpha_n C^{\frac{n}{n-1}} - \frac{n}{n-1} \Lambda_\epsilon} \int_{B_L} \frac{\epsilon^n}{(1 + c_n |x|^{\frac{n}{n-1}})^n} dx \\ &= e^{\alpha_n C^{\frac{n}{n-1}} - \frac{n}{n-1} \Lambda_\epsilon} (n-1) \epsilon^n \int_0^{c_n L^{\frac{n}{n-1}}} \frac{u^{n-2}}{(1+u)^n} du \\ &= e^{\alpha_n C^{\frac{n}{n-1}} - \frac{n}{n-1} \Lambda_\epsilon} (n-1) \epsilon^n \int_0^{c_n L^{\frac{n}{n-1}}} \frac{((u+1)-1)^{n-2}}{(1+u)^n} du \\ &= e^{\alpha_n C^{\frac{n}{n-1}} - \frac{n}{n-1} \Lambda_\epsilon} \epsilon^n (1 + O(L^{-\frac{n}{n-1}})) \\ &= \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \cdots + 1/(n-1)} \\ &\quad + O\left((L\epsilon)^n C^n \log L + L^{\frac{-n}{n-1}} + (L\epsilon)^n \log^n L\epsilon\right). \end{aligned}$$

Here, we used the fact

$$\sum_{k=0}^m \frac{(-1)^{m-k}}{m-k+1} C_m^k = \frac{1}{m+1}.$$

Then

$$\int_{B_{L\epsilon}} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx \geq \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)} + O\left((L\epsilon)^n C^n \log L + L^{\frac{-n}{n-1}} + (L\epsilon)^n \log^n L\epsilon\right).$$

Moreover, on $\mathbb{R}^n \setminus B_{L\epsilon}$ we have the estimate

$$\int_{\mathbb{R}^n \setminus B_{L\epsilon}} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx \geq \frac{\alpha_n^{n-1}}{(n-1)!} \int_{\mathbb{R}^n \setminus B_{L\epsilon}} \left| \frac{G(x)}{C^{\frac{1}{n-1}}} \right|^n dx,$$

and thus we get

$$\begin{aligned} \int_{\mathbb{R}^n} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx &\geq \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)} \\ &+ \frac{\alpha_n^{n-1}}{(n-1)!} \int_{\mathbb{R}^n \setminus B_{L\epsilon}} \left| \frac{G(x)}{C^{\frac{1}{n-1}}} \right|^n dx + O\left((L\epsilon)^n C^n \log L + L^{\frac{-n}{n-1}} + (L\epsilon)^n \log^n L\epsilon\right) \\ &= \frac{\omega_{n-1}}{n} e^{\alpha_n A + 1 + 1/2 + \dots + 1/(n-1)} \\ &+ \frac{\alpha_n^{n-1}}{(n-1)! C^{\frac{n}{n-1}}} \left[\int_{\mathbb{R}^n \setminus B_{L\epsilon}} |G(x)|^n dx + O\left((L\epsilon)^n C^{n+\frac{n}{n-1}} \log L + \frac{C^{\frac{n}{n-1}}}{L^{\frac{n}{n-1}}} + C^{\frac{n}{n-1}} (L\epsilon)^n \log^n L\epsilon\right) \right] \end{aligned} \quad (5.4)$$

We now set

$$L = -\log \epsilon; \quad (5.5)$$

then $L\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$. We then need to prove that there exists a $C = C(\epsilon)$ which solves equation (5.1). We set

$$f(t) = -\alpha_n t^{\frac{n}{n-1}} - (n-1)(1 + 1/2 + \dots + 1/(n-1)) + \alpha_n A + \log \frac{\omega_{n-1}}{n} - \log \epsilon^n + \phi,$$

Since

$$f\left(-\frac{2}{\alpha_n} \log \epsilon^n\right)^{\frac{n}{n-1}} = \log \epsilon^n + o(1) + \phi < 0$$

for ϵ small, and

$$f\left(-\frac{1}{2\alpha_n} \log \epsilon^n\right)^{\frac{n}{n-1}} = -\frac{1}{2} \log \epsilon^n + o(1) + \phi > 0$$

for ϵ small, f has a zero in $\left(-\frac{1}{2\alpha_n} \log \epsilon^n\right)^{\frac{n-1}{n}}, \left(-\frac{2}{\alpha_n} \log \epsilon^n\right)^{\frac{n-1}{n}}$. Thus, we defined C , and it satisfies

$$\alpha_n C^{\frac{n}{n-1}} = -\log \epsilon^n + O(1).$$

Therefore, as $\epsilon \rightarrow 0$, we have

$$\frac{\log L}{C^{\frac{n}{n-1}}} \rightarrow 0,$$

and then

$$(L\epsilon)^n C^{n+\frac{n}{n-1}} \log L + C^{\frac{n}{n-1}} L^{\frac{-n}{n-1}} + C^{\frac{n}{n-1}} (L\epsilon)^n \log^n L \epsilon \rightarrow 0.$$

Therefore, i), ii), iii) hold and we can conclude from (5.4) that for $\epsilon > 0$ sufficiently small

$$\int_{\mathbb{R}^n} \Phi(\alpha_n u_\epsilon^{\frac{n}{n-1}}) dx > \frac{\omega_{n-1}}{n} e^{\alpha_n A+1+1/2+\dots+1/(n-1)}.$$

6 The test function 2

In this section we construct, for $n > 2$, functions u_ϵ such that

$$\int_{\mathbb{R}^n} \Phi(\alpha_n (\frac{u_\epsilon}{\|u_\epsilon\|_{H^{1,n}}})^{\frac{n}{n-1}}) dx > \frac{\alpha_n^{n-1}}{(n-1)!},$$

for $\epsilon > 0$ sufficiently small.

Let $\epsilon^n = e^{-\alpha_n c^{\frac{n}{n-1}}}$, and

$$u_\epsilon = \begin{cases} c & |x| < L\epsilon \\ \frac{-n \log \frac{x}{L}}{\alpha_n c^{\frac{n}{n-1}}} & L\epsilon \leq |x| \leq L \\ 0 & L \leq |x|, \end{cases}$$

where L is a function of ϵ which will be defined later.

We have

$$\int_{\mathbb{R}^n} |\nabla u_\epsilon|^n = 1,$$

and

$$\int_{\mathbb{R}^n} u_\epsilon^n dx = \frac{\omega_{n-1}}{n} c^n (L\epsilon)^n + \frac{\omega_{n-1} n^n L^n}{\alpha_n^n c^{\frac{n}{n-1}}} \int_\epsilon^1 r^{n-1} \log^n r dr.$$

Then

$$\begin{aligned} \int_{\mathbb{R}^n} \Phi(\alpha_n (\frac{u_\epsilon}{\|u_\epsilon\|_{H^{1,n}}})^{\frac{n}{n-1}}) dx &\geq \frac{\alpha_n^{n-1}}{(n-1)!} \frac{\int_{\mathbb{R}^n} u_\epsilon^n dx}{1 + \int_{\mathbb{R}^n} u_\epsilon^n dx} + \frac{\alpha_n^n}{n!} \frac{\int_{\mathbb{R}^n \setminus B_{L\epsilon}} u_\epsilon^{\frac{n^2}{n-1}}}{(1 + \int_{\mathbb{R}^n} u_\epsilon^n dx)^{\frac{n}{n-1}}} dx \\ &= \frac{\alpha_n^{n-1}}{(n-1)!} - \frac{\alpha_n^{n-1}}{(n-1)!} \frac{1}{1 + \frac{\omega_{n-1}}{n} c^n (L\epsilon)^n + \frac{\omega_{n-1} n^n L^n}{\alpha_n^n c^{\frac{n}{n-1}}} \int_\epsilon^1 r^{n-1} \log^n r dr} \\ &\quad + \frac{\alpha_n^n}{n!} \frac{\omega_{n-1} L^n / c^{\frac{n^2}{(n-1)^2}} (\frac{n}{\alpha_n})^{\frac{n^2}{n-1}} \int_\epsilon^1 r^{n-1} \log^{\frac{n^2}{n-1}} r}{\left(1 + \frac{\omega_{n-1}}{n} c^n (L\epsilon)^n + \frac{\omega_{n-1} n^n L^n}{\alpha_n^n c^{\frac{n}{n-1}}} \int_\epsilon^1 r^{n-1} \log^n r dr\right)^{\frac{n}{n-1}}} \end{aligned}$$

We now ask that L satisfies

$$\frac{c^{\frac{n}{n-1}}}{L^n} \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0. \quad (6.1)$$

Then, for sufficiently small ϵ , we have

$$\begin{aligned}
& -\frac{\alpha_n^{n-1}}{(n-1)!} \frac{1}{1 + \frac{\omega_{n-1}}{n} c^n (L\epsilon)^n + \frac{\omega_{n-1} n^n L^n}{\alpha_n^n c^{\frac{n}{n-1}}} \int_\epsilon^1 r^{n-1} \log^n r dr} + \\
& + \frac{\alpha^n}{n!} \frac{\omega_{n-1} L^n / c^{\frac{n^2}{(n-1)^2}} (\frac{n}{\alpha_n})^{\frac{n^2}{n-1}} \int_\epsilon^1 r^{n-1} \log^{\frac{n^2}{n-1}} r}{\left(1 + \frac{\omega_{n-1}}{n} c^n (L\epsilon)^n + \frac{\omega_{n-1} n^n L^n}{\alpha_n^n c^{\frac{n}{n-1}}} \int_\epsilon^1 r^{n-1} \log^n r dr\right)^{\frac{n}{n-1}}} \\
& \geq B_1 L^{n - \frac{n^2}{n-1}} - B_2 \frac{c^{\frac{n}{n-1}}}{L^n} \\
& = \frac{c^{\frac{n}{n-1}}}{L^n} (B_1 \frac{L^{2n - \frac{n^2}{n-1}}}{c^{\frac{n}{n-1}}} - B_2) \\
& = \frac{c^{\frac{n}{n-1}}}{L^n} (B_1 \frac{L^{\frac{n}{n-1}(n-2)}}{c^{\frac{n}{n-1}}} - B_2) ,
\end{aligned}$$

where B_1, B_2 are positive constants.

When $n > 2$, we may choose $L = b c^{\frac{1}{n-2}}$; then, for b sufficiently large, we have

$$B_1 \frac{L^{\frac{n}{n-1}(n-2)}}{c^{\frac{n}{n-1}}} - B_2 = B_1 b^{\frac{n}{n-1}(n-2)} - B_2 > 0 ,$$

and (6.1) holds. Thus, we have proved that for $\epsilon > 0$ sufficiently small

$$\int_{\mathbb{R}^n} \Phi(\alpha_n (\frac{u_\epsilon}{\|u_\epsilon\|_{H^{1,n}(\mathbb{R}^n)}})^{\frac{n}{n-1}}) dx > \frac{\alpha_n^{n-1}}{(n-1)!} .$$

References

- [A-T] S. Adachi and K. Tanaka: Trudinger type inequalities in $\mathbb{R}^n$ and their best exponents. Proc. AMS, **128**: 2051-2057, 1999.
- [A] D. R. Adams: A sharp inequality of J. Moser for higher order derivatives. Annals of Math., **128**: 385-398, 1988.
- [A-D] Adimurthi and O. Druet, Blow up analysis in dimension 2 and a sharp form of Trudinger Moser inequality, Comm. PDE **29** No 1-2, (2004), 295-322.
- [A-M] Adimurthi and M. Struwe: Global Compactness properties of semilinear elliptic equations with critical exponential growth. J. Funct. Anal., **175**, no 1:125-167, 2000.
- [C-C] L. Carleson and S. Y. A. Chang: on the existence of an extremal function for an inequality of J. Moser. Bull. Sc. Math., **110**, 113-127, 1986.
- [d-d-R] D.G. de Figueiredo, J.M. do O', B. Ruf: On a inequality by N. Trudinger and J. Moser and related elliptic equations. Comm. Pure. Appl. Math., **55**, 135-152, 2002.
- [D] E. DiBenedetto: $C^{1,\alpha}$ local regularity of weak solution of degenerate elliptic equations. Nonlinear Analysis. **7** no.8, 827-850, 1983.

- [F] M. Flucher: Extremal functions for Trudinger-Moser inequality in 2 dimensions. *Comm. Math. Helv.*, **67**,471-497, 1992.
- [Fo] L. Fontana: Sharp borderline Sobolev inequalities on compact Riemannian manifolds. *Comm. Math. Helv.*, **68**,415- 454, 1993.
- [K-L] S. Kichenassamy and L. Veron: Singular solutions of the p-laplace equation. *Math. Ann.*, **275**,599-615, 1986.
- [L1] Y. Li: Moser-Trudinger inequality on manifold of dimension two. *J. Partial Differential Equations* **14**, no. 2, 163–192, 2001.
- [L2] Y. Li: The extremal functions for Moser-Trudinger inequality on compact Riemannian manifolds, To appear in *Sci. Chinese, series A*.
- [L3] Y. Li: Remarks on the Extremal Functions for the Moser-Trudinger Inequalities. To appear in *Acta Mathematica Sinica*.
- [L-L] Y. Li and P. Liu: A Moser-Trudinger inequality on the boundary of a Riemann Surface. *Math. Z.*, **250**, no.2, 363 - 386, 2005.
- [Lin] K.C. Lin: Extremal functions for Moser's inequality. *Trans. Amer. Math. Soc.*, **348**,2663-2671, 1996.
- [M] J. Moser: A sharp form of an Inequality by N.Trudinger. *Ind. Univ. Math. J.*, **20**,1077-1091, 1971.
- [P] S. I. Pohozaev: The Sobolev embedding in the case $p_1 = n$. *Proceedings of the Technical Scientific Conference on Advances of Scientific Research 1964-1965. Mathematics Section*, 158-170, Moskov. Energet. Inst., Moscow, 1965
- [R] B. Ruf: A sharp Trudinger-Moser type inequality for unbounded domains in $\mathbb{R}^2$. *J. Funct. Anal.* **219**, no. 2, 340–367, 2005.
- [T] P. Tolksdorf: Regularity for a more general class of quasilinear elliptic equations. *J.D.E.*, **51**:126-150 1984.
- [S] J. Serrin: Local behavior of solutions of quasilinear equations. *Acta. Math.*, **111**,248-302, 1964.
- [St] M. Struwe: Positive solution of critical semilinear elliptic equations on non-contractible planar domain. *J. Eur. Math. Soc.*, **2**(4):329–388, 2000.
- [Tr] N. S. Trudinger: On embedding into Orlicz space and some applications, *J. Math. Mech.* **17**:473-484, 1967.

Yuxiang Li

ICTP, Mathematics Section, Strada Costiera 11, I-34014 Trieste, Italy

E-mail address: liy@ictp.it

Bernhard Ruf

Dipartimento di Matematica, Università di Milano, via Saldini 50, 20133 Milan, Italy

E-mail address: Bernhard.Ruf@mat.unimi.it